

KRETOV, A. Ye.; BORODAVKO, N. D.

N, N-di (β -cyanoethyl)cyanamide and its reactions. Zhur. ob.
khim. 33 no. 5:1536-1539 My '63. (MIRA 16:6)

(Cyanamide)

KRETOV, A.Yo.; BESPALYZ, A.S.

Derivatives of thiazolidine. Zhur.ob.khim. 33 no.6:1878-1882
Je '63. (MIRA 16:7)
(Thiazolidine)

KRETOV, A.Ye.; ABRAZHANOVA, Ye.A.; ZLOTCHENKO, S.I.; KUKHAR', V.P.

Arene sulfamido ketones. Zhur.ob.khim. 33 no.7:2355-2357 J1
'63. (MIRA 16:8)
(Acetophenone) (Sulfamide)

KRETOV, A.Ye.; MOMSENKO, A.P.

Reactions of cyanamide with aliphatic acid anhydrides. Zhur.ob.
khim. 33 no.10:3325-3328 0 '63. (MIRA 16:11)

1. Dnepropetrovskiy khimiko-tehnologicheskii institut.

KRETOV, A.Ye.; STERINA, Ye.Z.

Acenaphtholylacrylic acids and their derivatives. Zhur. prikl.
khim. 36 no.5:1154-1157 My '63. (MIRA 16:8)

(Naphthaleneacrylic acid)

KOZOPOLYANSKIY, N.S.; KRETOV, A.Ye; OKHRAMOVICH, A.Ye.; ILYASH, I.I.

Use of fluorene-9,9-dipropionic acid for modification of
polyester resins. Plast. massy no.11:14-15 '63. (MIRA 16:12)

KRETOV, A.Ye.; BESPALYY, A.S.

Derivatives of thiazolidine. Part 2. Zhur.ob.khim. 33 no.10:
3323-3325 0 '63. (MIRA 16:11)

1. Dnepropetrovskiy khimiko-tekhnologicheskii institut imeni
F.E.Dzerzhinskogo.

OKHRAMOVICH, A.Ye.; KRETOV, A.Ye.

Preparation of polyesters by the condensation of fluorene-9,
9-dipropionic acid with 1,4-butylen glycol and 1,1-dihydroxyethyl
ester. Zhur. prikl. khim. 36 no.12.2775-2779 D'63.

(MIRA 17:2)

1. Dnepropetrovskiy khimiko-tekhnologicheskii institut.

KRETOV, A. Ye.; TIKHONOVA, G.V.

Reactions of diglymidamide with acetaldehyde and chloral.
Zhur. ob. khim. 34 no.7:2428-2430 JI '64 (MIRA 17:8)

1. Dnepropetrovskiy khimiko-tekhnologicheskii Institut.

KRETOV, A.Ye.; BESPALYY, A.S.

Derivatives of naphthothiazinidine. Zhur. ob. khim. 34 no. 3:
999-1001 Mr '64. (MIRA 17:6)

1. Dnepropetrovskiy khimiko-tehnologicheskii institut.

KROTOV, A. Ye.; BES-PALYY, A. S.; POLITUN, N. N.

Thiophenolsulfonic acids and their derivatives. *Zhur. ob.*
Khim. 34 no.6/2066-2068 Js '64. (NIR: 1711)

1. Dnepropetrovskiy khimiko-tekhnologicheskii institut.

OKHRAMOVICH, A.Ye.; KRETOV, A.Ye.

Esters of fluorene-9,9-dipropionic acid. Zhur.prikl.khim. 37 no.1:
220-223 Ja '64. (MIRA 17:2)

1. Dnepropetrovskiy khimiko-tehnologicheskii institut.

GOLOS, I.O.; KUPCHEN, A.S.; POLOV, R.A.

Synthesis of thiazolidine-5-acetic acid derivatives. *Trav. ca.*
khim. 34 no.9:3063-3066 1964. (MIRA 17:10)

I. Maspropetrovskiy khimiko-tekhnologicheskoy institut.

KRETOV, A.Ye.; BESFALYY, A.S.

Thiazolidine derivatives. Part 3. Zhur. ob. khim. 34 no.10:3365-3367
O '64. (MIRA 17:11)

1. Dnepropetrovskiy khimiko-tehnologicheskii institut imeni F.E.
Dzerzhinskogo.

KHELOV, A.Ye.; DAVYDOV, A.V.

New method of synthesizing guanamines containing fluoroalkyl radicals. Zhur. ob. khim. 35 no.4:746-748 Ap '65.

(MIRA 18:5)

1. Dnepropetrovskiy khimiko-tehnologicheskii institut.

KRETOV, A.Ye.; OKHRAMOVICH, A.Ye.

Preparation of di- and tri-(β -cyanoethyl)-indene and their derivatives. Zhur.prikl.khim. 37 no.7:1617-1619 J1 '64.

(MIRA 18:4)

1. Dnepropetrovskiy khimiko-tekhnologicheskii institut.

LEVIN, A.I. [Borisov]; LEVIN, A.V.

Study of the reaction of fluorocarboxylic acids with tin(II).
Zhur. ob. khim. 35 no.7:1156-1158 J1 '65. (1965)

1. Inopropetrovskiy khimiko-tekhnologicheskoy institut.

ERSTOV, A.Ye. [deceased]; ABRAZHANOVA, Ye.A.; KUKHAR', V.P.

Oximes of alkoxy- and aroxy-cyclohexanones. Zhur. org. khim. 1 no.6:
1021-1022 Jo '65. (MIRA 18:7)

ZLATIN, L.I.; KRETOV, B.K.

Automatic opening of the gates of coke ramps. Koks i khim. no.1:
41-45 '56. (MIRA 9:5)

1. Kemerovskiy koksokhimicheskiy zavod.
(Coke industry--Equipment and supplies)

KRETOV B.K.

AUTHOR: Zlatin, L.I. and Kretov, B.K.

68-12-22/25

TITLE: Mechanization of Loading Ammonium Sulphate in Box
Cars (Kompleksnaya mekhanizatsiya pogruzki sul'fata
ammoniya v krytyye vagony)

PERIODICAL: Koks i Khimiya, 1957, No.12, pp. 50 - 52 (USSR)

ABSTRACT: Mechanization of loading ammonium sulphate in covered
wagons, organised on the Kemerov Coke Oven Works, is described
and illustrated. There are 3 figures.

ASSOCIATION: Kemerovo ~~chemical plant~~ zavod (Kemerovskiy koksokhimicheskiy
zavod)

AVAILABLE: Library of Congress

Card 1/1

ZLATIN, L.I.; KRETOV, B.K.; PANENKO, F.M.

Use of self-sealing doors in pitch coke ovens. Koks i khim. no.4:51
'60. (MIRA 13:6)

1. Kemerovskiy koksokhimicheskiy zavod.
(Kemerovo--Coke ovens)

KRETOV, G., inzh.

Strictly speaking, it is correct. But is it really? Grazhd.
av. 20 no.3:10-11 Mr '63. (MIRA 16:4)

(Aeronautics, Commercial)

PAVLOVSKIY, V.Ya.; TSILEVICH, I.Z.; FRADIN, M.P.; LECHTAFONICH, P.D.;
SHAPIRO, Yu.A.; GRIGOR'YEVA, M.G.; RABNO-LINA, Ye.T.; KERTOVA, G.V.

Rolling mill rolls of hypereutectoid chromium-vanadium 90 KhF steel.
Metallurg 10 no.7:40 J1 '65. (MIRA 12:7)

1. Metallurgicheskiy zavod "Azovstal'".

KRETOV, M.

1. 1964 1965 1966 1967 1968 1969 1970 1971 1972 1973 1974 1975 1976 1977 1978 1979 1980 1981 1982 1983 1984 1985 1986 1987 1988 1989 1990 1991 1992 1993 1994 1995 1996 1997 1998 1999 2000 2001 2002 2003 2004 2005 2006 2007 2008 2009 2010 2011 2012 2013 2014 2015 2016 2017 2018 2019 2020 2021 2022 2023 2024 2025 2026 2027 2028 2029 2030 2031 2032 2033 2034 2035 2036 2037 2038 2039 2040 2041 2042 2043 2044 2045 2046 2047 2048 2049 2050 2051 2052 2053 2054 2055 2056 2057 2058 2059 2060 2061 2062 2063 2064 2065 2066 2067 2068 2069 2070 2071 2072 2073 2074 2075 2076 2077 2078 2079 2080 2081 2082 2083 2084 2085 2086 2087 2088 2089 2090 2091 2092 2093 2094 2095 2096 2097 2098 2099 2100 2101 2102 2103 2104 2105 2106 2107 2108 2109 2110 2111 2112 2113 2114 2115 2116 2117 2118 2119 2120 2121 2122 2123 2124 2125 2126 2127 2128 2129 2130 2131 2132 2133 2134 2135 2136 2137 2138 2139 2140 2141 2142 2143 2144 2145 2146 2147 2148 2149 2150 2151 2152 2153 2154 2155 2156 2157 2158 2159 2160 2161 2162 2163 2164 2165 2166 2167 2168 2169 2170 2171 2172 2173 2174 2175 2176 2177 2178 2179 2180 2181 2182 2183 2184 2185 2186 2187 2188 2189 2190 2191 2192 2193 2194 2195 2196 2197 2198 2199 2200 2201 2202 2203 2204 2205 2206 2207 2208 2209 2210 2211 2212 2213 2214 2215 2216 2217 2218 2219 2220 2221 2222 2223 2224 2225 2226 2227 2228 2229 2230 2231 2232 2233 2234 2235 2236 2237 2238 2239 2240 2241 2242 2243 2244 2245 2246 2247 2248 2249 2250 2251 2252 2253 2254 2255 2256 2257 2258 2259 2260 2261 2262 2263 2264 2265 2266 2267 2268 2269 2270 2271 2272 2273 2274 2275 2276 2277 2278 2279 2280 2281 2282 2283 2284 2285 2286 2287 2288 2289 2290 2291 2292 2293 2294 2295 2296 2297 2298 2299 2300 2301 2302 2303 2304 2305 2306 2307 2308 2309 2310 2311 2312 2313 2314 2315 2316 2317 2318 2319 2320 2321 2322 2323 2324 2325 2326 2327 2328 2329 2330 2331 2332 2333 2334 2335 2336 2337 2338 2339 2340 2341 2342 2343 2344 2345 2346 2347 2348 2349 2350 2351 2352 2353 2354 2355 2356 2357 2358 2359 2360 2361 2362 2363 2364 2365 2366 2367 2368 2369 2370 2371 2372

• • • • •

(Switzerland--D.)

(111)

KRETOV, I.T.

Drying brewer's waste. Izv. vys. ucheb. zav.; pishch. tekhn.
no.4:124-127 '61. (MIRA 14:8)

1. Leningradskiy tekhnologicheskii institut pishchevoy promyshlennosti,
kafedra oborudovaniya pishchevykh predpriyatiy.
(Brewing industry--By-products)

KRETOV, I.T.

Determining the coefficient of moisture transfer in brewing grains. Izv.vys.ucheb.zav.; pishch.tekh. 2:138-143 '62.

(MIRA 15:5)

1. Voronezhskiy tekhnologicheskii institut, kafedra oborudovaniya pishchevykh predpriyatiy.

(Brewing industry) (Grain--Drying)

PP(1)

JOV/47-59-3-24/53

AUTHOR: Kretov N.A.

TITLE: The Regeneration of Permanent Magnets

PERIODICAL: Fizika v shkole, 1959, Nr 3, p 75 (USSR)

ABSTRACT: The author describes a method to regenerate weak permanent magnets. The magnet is placed within a 12-volt coil taken from a demountable school transformer. The poles of the magnet are closed with an iron armature. For a linear magnet, the magnetic circuit can be established with the core parts of the transformer. The coil is switched to a 127 or 220 volt alternating current circuit through a copper conductor of 0.15 to 0.2 mm. Upon contact, the thin conductor burns out, having time to let current pass in one direction only. The polarity of the regenerated magnets can be checked with a magnetic needle.

Card 1/2

SOV/47-59-3-24/53

The Regeneration of Permanent Magnets

Berezhnyakovskaya srednyaya shkola Usmanskogo rayona
Lipetskoy obl. (Berezhnyakovskaya Secondary School of the
Usman' Rayon, Lipetsk Oblast)

Card 2/2

RUKHLYADEVA, A.P.; POLYGALINA, G.V.; BAULINA, E.A.; KRETOV, V.F.

Automatic method for determining the concentration of grain and potato mash. Ferm. i spirt. prom. 30 no.3:25-29 '64.

(MIRA 18:2)

1. Vsesoyuznyy nauchno-issledovatel'skiy institut fermentnoy i spirtovoy promyshlennosti (for Rukhlyadeva, Polygalina).
2. Vsesoyuznyy nauchno-issledovatel'skiy eksperimental'no-konstruktorskiy institut prodovol'stvennogo mashinostroyeniya (for Baulina, Kretov).

ODINOKOV, S.D., kand.tekhn.nauk; SHABALINA, V.I., mladshiy nauchnyy
sotrudnik; SIROTKINA, O.V., starshiy tekhnika; KRETOVA, L.V.,
starshiy tekhnika; VDOVENKO, Z.I., red.izd-va; TEMKINA, Ye.L.,
tekhn.red.

[Album of charts, designs of equipment, tools, and devices for
erecting asbestos cement building roofs] Al'bom tekhnologi-
cheskikh skhem, chertezhei oborudovaniya, instrumentov i prispo-
soblenii dlia ustroystva asbestotsementnykh krovel' zdaniy.
Moskva, Gos.izd-vo po stroit., arkhitekt. i stroit.materialam, 1960.
42 p. (MIRA 14:3)

1. Akademiya stroitel'stva i arkhitektury SSSR. Institut orga-
nizatsii, mekhanizatsii i tekhnicheskoy pomoshchi stroitel'stvu.
2. Laboratoriya krovel'nykh i otdelochnykh rabot Nauchno-issle-
dovatel'skogo instituta organizatsii, mekhanizatsii i tekhnicheskoy
pomoshchi stroitel'stvu Akademii stroitel'stva i arkhitektury SSSR
(for Odinokov, Shabalina, Sirotkina, Krotova).
(Asbestos cement) (Roofing)

NOSKOV, S.K., kand.tekhn.nauk; ODINOKOV, S.D., kand.tekhn.nauk; SIROTKINA, O.V., starshiy tekhnik; KRETOVA, L.V., starshiy tekhnik. Prinimala uchastiye SHABALINA, V.I., mladshiy nauchnyy sotrudnik. SKVORTSOVA, I.P., red.izd-va; TEMKINA, Ye.L., tekhn.red.

[Album of technological schemes and drawings of the equipment, instruments, and devices to be used in covering roofs with rolled materials] Al'bom tekhnologicheskikh skhem i chertezhei oborudovaniia, instrumentov i prispособlenii dlia ustroistva krovvel' iz rulonnykh materialov. Moskva, Gos.izd-vo lit-ry po stroit., arkhitekt. i stroit.materialam, 1960. 48 p. (MIRA 13:6)

1. Akademiya stroitel'stva i arkhitektury SSSR. Institut organizatsii, mekhanizatsii i tekhnicheskoy pomoshchi stroitel'stvu. (Roofing--Equipment and supplies)

KRETOV, L.Ve.; MOMSENKO, A.P.

Reactions of cyanamide with aliphatic acid anhydrides. Part 1.
Zhur.ob.khim. 33 no.2:397-399 F '63. (MIRA 1642)

1. Dnepropetrovskiy khimiko-tekhnologicheskii institut.
(Cyanamide) (Acids, Fatty) (Anhydrides)

SULIMOV, Filaret Ivanovich; GORBACHEV, Sergey Mikhaylovich;
KRETOV, Pavel Yevseyevich; LIOGEN'KIY, German L'vovich;
VELISHCHANSKIY, V.M., red.; YELCHINA, L.A., red.izd-va;
KAZANSKAYA, L.I., tekhn.red.

[Reorganization problems and forest management in Vologda
Province] Voprosy reorganizatsii i lesnoe khoziaistvo
Vologodskoi oblasti. Moskva, Goslesbumizdat, 1963. 74 p.
(MIRA 17:3)

KRETOV, V.P., inzh.; KIRPICHNIKOV, Yu.A., inzh.

Automatic lighting control. Prom. energ. 19 no.11:22-23 N 164.
(MIR: 18:1)

KRETOVA, N. F.

112-6-11867

Translation from: Referativnyy zhurnal, Elektrotehnika, 1957, No. 5, p. 15 (USSR)

AUTHOR: Voronkov, G.N., Zvyagil'skiy, A.A., and Kretova, N.F.

TITLE: High-Voltage Porcelain of Better Electromechanical Properties from Boron-Containing Raw Material (Vysokovol'tnyy farfor s povyshennymi elektromekhanichesкими свойствами na osnove borosoderzhashchego syr'ya)

PERIODICAL: Tr. Gos. issled. elektrokeram. in-ta, 1956, Nr. 5, pp. 5-16

ABSTRACT: As it was necessary to improve the mechanical and electrical characteristics of porcelain a new type of porcelain was developed in GIEKI on the basis of a boron-containing (ascharit) ore, alumina, clay materials and a small amount of alkali-earth compounds. No quartz or feldspar was introduced. The use of ascharite ore ($2\text{MgO} \cdot \text{B}_2\text{O}_3 \cdot \text{H}_2\text{O}$) as a fusing agent, instead of CaCO_3 or BaCO_3 , and also the introduction of commercial Al_2O_3 with an increased content of kaolin insured the close-packed structure of porcelain, in which the crystals of mullite formed a felt-like lattice and were uniformly distributed in the vitreous phase. There is a negligible amount of free sections of glass in the ascharite porcelain, but there are finely grained clusters of α -alumina. As the ascharite porcelain has a lower coefficient of linear expansion (3.9×10^{-6}) than the ordinary feldspar porcelain (6×10^{-6}), two new glazes (white and brown) were developed having less alkali oxide content. Due to

Card 1/2

112-6-11867

Translation from: Referativnyy zhurnal, Elektrotehnika, 1957, Nr. 1, p. 13 (USSR)

the more uniform structure and other factors the ascharite porcelain has almost double mechanical strength as compared to the feldspar porcelain. Nonalkaline vitreous phase insures higher values of volume electrical resistivity and electric strength, and lower values of the dielectric loss angle. Preparatory procedures and the manufacture of insulators can follow regular methods of the electrical porcelain manufacture. The only additional operation is the introduction of sinter into the mass of ascharite porcelain. Optimum firing temperature 1310 - 1330°C. Ascharite and feldspar insulators can be fired jointly, but the sintering interval of the ascharite units is shorter than that of the ordinary electrical porcelain (30-40° against 60-80°C). Thermographic and chemical investigations of the ascharite ore have shown that for electrical porcelain purposes it should have at least 22% B_2O_3 and 23% MgO . The density of ascharite ore should be at least 2.67 g/cm³, the firing loss should not be over 18%. Bibliography: 6 titles.

H.V.N.

Card 2/2

Dissertation: "Investigation of the Resistance to Rolling of a Tractor with Pneumatic Tires"
Cand Tech Sci, All-Union Sci Res Inst of Mechanization of Agriculture, Moscow, 1953.
(Referativnyy Zhurnal--znanie, Moscow, Apr 54)

SO: SUK 243, 19 Oct 1954

KRETOVA, G.

KRETOVA, G. "The Kamennaya Steppe. In the natural reservation", (Outline), Lit. Voronezh, 1946, No. 3, p. 159-200.

SO: U-3042, 11 March 53, Letopis: 'Zhurnal 'nykh Statoy, No.7 1949).

KRETOVA, O.

Kretova, O. "The rocky steppe," [On the work of the Farming
Institute of the Central Chernozem Zone imeni Dokuchaevskiy],
Oktyabr', 1949, No. 3, p. 102-34

SO: U-3566, 15 March 53, (Letopis 'Zhurnal 'nykh Statay, No. 14, 1949).

KRETOVA, O., pisatel'; BULAVIN, M., pisatel'; GLUKHOV, A., kand.ekon.nauk;
MITROSHIN, S., kand.istoricheskikh nauk; PLOTNIKOV, A., vrach;
MOREV, M., zhurnal'st; PRUDKOVSKIY, P.N., red.; VOROTNIKOVA, R.V.,
red.; SZERADZSKAYA, P.G., tekhn.red.

[From impoverishment to prosperity; past and present conditions of the
villages of Novo-Zhivotinnoye and Mokhovatka, Berezov District, Voro-
nesh Province] Ot oskudeniya k protsvetaniyu; proshloe i nastoiashchee
sel Novo-Zhivotinnogo i Mokhovatki Berezovskogo raiona Voroneshskoi
oblasti. Voroneshskoe knizhnoe izd-vo, 1958. 77 p. (MIRA 12:3)

1. Zaveduyushchiy Novo-Zhivotinnovskoy uchastkovoy bol'nitsy (for
Plotnikov).

(Voronesh Province--Villages)

KRETOVA, Ol'ga Kapitonovna; PRUDKOVSKIY, P.N., red.; SERADZSKAYA, P.O.,
tekhn.red.

[We who live near Voronezh; a sketch] Pod Voronezhem u nas;
ocherk. Voronezh, Voronezhskoe knizhnoe izd-vo, 1959. 27 p.
(MIRA 14:1)

(Manukovskii, Nikolai Fedorovich)

KRETOVA, Ol'ga Kapitonovna; DROKHANOVA, Ye.N., red.; YELAGIN, A.S.,
tekh. red.

[Nikolai Mamukovskii's "universities."] Universitety Nikolaia
Mamukovskogo. Moskva, Izd-vo "Sovetskaia Rossiia, 1961. 124 p.
(MIRA 15:3)

(Mamukovskii, Nikolai Fedorovich)

KRETOVA, T.S.; SMIRNOVA, N.P., redakter; MAKHOVA, N.N., tekhnicheskii redakter.

[The teacher's preparation for geography lessons in class 5] Podgotovka uchitelia k urokam geografii v V klasse. Moskva, Gos.uchebno-pedagog. izd-vo Ministerstva prosveshcheniia RSFSR, 1954. 47 p.
(Physical geography--Study and teaching) (MIRA 8:5)

1. KRETOVA, V. S.

2. SSSR (600)

4. Geese

7. My work practice.
Ptitsevodstvo No. 6, 1952

9. Monthly List of Russian Accessions, Library of Congress, February 1953, Unclassified.

KRETOVICH, V. L.

[illegible]

830 524 METALLURGICAL LITERATURE CLASSIFICATION

The distribution of sugar and nitrogen compounds in
 wheat grain. V. Kretovich. *Sov. Inst. Cereal Research*
 (Moscow) 13, 70-3(1934). Sucrose is present not only
 in the kernel, but in much greater amts. also in the endo-
 sperm. The sugar concn. in the outer layers of the endo-
 sperm is 4.8 times more than that of the inner layer.
 In the aleurone layer, there is no sugar. Of the proteins,
 gliadin, the chief component of gluten, is lacking. The
 main part of the proteins in the aleurone layer consists of
 gluten and albumin. The expts. were performed with
 wheat grain from the harvest of 1911 (Crimea). H. C.

Biochemical changes in wheat grain under the action of high temperatures. V. L. Koryukh and K. N. Riasantseva. *Compt. rend. acad. sci. U. R. S. S. (N. S. I.)*, 3, 400-12(1933); cf. *C. A.*, 29, 4908. Samples of gliadin grains were heated in a chamber and the wdy. of gliadin in H_2O , hydration of gluten and the activities of catalase and diastase measured. Denaturation of gliadin was increased on heating to 90-130°. The hydration of gluten was markedly decreased. Catalase activity is sharply cut on heating to 90° even with quite dry grain. The diastase activity increased markedly at 90-95°, and only relatively high temp., 105°, caused a decrease in H_2O , but in phosphate-citrate buffer it is resistant to even these temps. Control of grain drying and hot-conditioning could be based upon detn. of catalase activity. C. P. P. J.

ASB-3LA METALLURGICAL LITERATURE CLASSIFICATION


111 AND 112 ORDERS		PROCESSING AND PROPERTIES INDEX	
<p>The relation between gliadin and glutenin in the wheat grain. V. L. Kretovich and E. Ryazantsev. <i>Trudy Lab. Tsvetnykh Belkh Polkovogo Otmennogo Organismo</i> 8, 82 4 (1935); <i>Chem. Zentr.</i> 1938, II, 3175; cf. C. A. 30, 1445⁹.—The amt. of alc.-sol. N contained in the flour varies with the thickness of the peripheral layers of the grain. Therefore, in testing the theory regarding the relation between the ratio of gliadin to glutenin on the one hand and the technological properties of the grain on the other, it is necessary that conditions and ratios in the bran be considered and not those in the flour, since only the content in the bran and not the total N in the grain is the chief factor detg. the plasticity and taste of pastries and bread. It was also shown that alc. solns. of gliadin very readily penetrate the cell walls of the intact grain, so that the degree of milling of the flour is without influence on the ratio of alc.-sol. N to total N. W. A. Morse</p>			
<p>ASAC-11A METALLURGICAL LITERATURE CLASSIFICATION</p>			
111 AND 112 ORDERS		111 AND 112 ORDERS	

17

Enzymes as a factor in the quality of grain and flour
V. L. Kretovskh. *Sprskhi Khim. S.* 1041-51 (1936). A
K. H. Rathmann

12

ASB-51A METALLURGICAL LITERATURE CLASSIFICATION



118

Biochemical changes in the grain of wheat damaged by the wheat-bug. V. Kretovich and R. Tokareva. *Biochimica J.* 387-407 (1978). The proteins of the damaged grain become very sol. in water as well as in 60% alc.; diastatic activity increases, and the acidity is also somewhat higher. The glutathione content is the same. The gluten from damaged grains shows a lower viscosity and specific rotation, and an increased S content. The damage to the grain is done only at the point bitten, and is not transmitted to the entire grain. H. Cohen

BIOCHEMICAL LABORATORY OF THE ALL-UNION GRAIN INSTITUTE

ADD. 3.1.1. BIOCHEMICAL LITERATURE CLASSIFICATION

KRETOVICH, V. L.

"The Biochemistry of Grain in Storage" A. I. Smirnov, and V. L. Kretovich,
Sbornik Akad Nauk SSSR, Presidentu Adad Nauk SSSR Komarovu 1939, pp 720-5;
Khim Referat Zhur, 1940, No 12, pp 31 (SEE: Inst. Insect/Fungi. in Ya. V.
Samoylov)

SO: U-237/49, 8 April 1949

17

No. 1

118

Biochemical residues of frost-damaged grain. V. Kiselevich and P. Tokareva. *Biohimiya* 4, 585-590 (1959).
Frost-damaged grain is characterized by an increased diastatic activity and acidity, and an inelasticity of gluten. Besides conditioning at 40° for 40 min., addn. of lactic acid improves the quality of the bread. H. Cohen

Lab. Biochem. All Union Res. inst. Moscow

ASD 34.4 METALLURGICAL LITERATURE CLASSIFICATION

12

CA

12

Biochemical changes in the grains of wheat damaged by the wheat bug H. Y. L. Kulyukh and P. P. Tokareva. *Biokhimiya* 4, 636-47 (1969); cf. *U. S. J.* 33, 683. The addn. of dry enzyme preps. from the damaged grains to normal wheat flour destroys the gluten. These enzymes are not activated by cysteine nor inhibited by bromates. Normal gluten was obtained from a wheat sample contg. 32% damaged grain by steaming for 1.5-3 min., or by heating (after the moisture content had been raised 20%) 30-45 min., rapidly raising the temp. from 50° to 80°.

H. Proskov

Biochemical Lab., All Union Inst. of Grain, Moscow

ASB 51.6 METALLURGICAL LITERATURE CLASSIFICATION

Effect of relative air humidity and anaerobiosis on freshly harvested grain of wheat. V. L. Krasovitsky, A. I. Sokolova, and E. N. Uschakova (Compt. rend. Acad. Sci. U.R.S.S., 1940, 23, 487—490).—Germinating power and energy of stored grain fall, but catalase and tyrosinase activity, free fatty acid, and non-protein-N contents rise, with increasing R.H. Optimum R.H. is 0—30%. After-ripening is not affected by absence of O_2 , but is retarded by storage in CO_2 . R. L. E.

12

CH

THE STABLE MOISTURE CONTENT OF GRAIN AND ITS EFFECT ON THE LIPASE ACTION. V. L. Krasovich, A. I. Sokolova and P. N. Ushakova. *Compt. rend. acad. sci. U. R. S. S.* 27, 1701-4 (1940) (in German).—For detn. of the acidity

No. 7
of the ether ext. (which is a measure of fat change and, therefore, of lipase action) 5 g. of ground wheat was extd. for 1 hr. with dry ether in a Soxhlet app. to 60 ml., and the ext. obtained was titrated with 0.01 N lye with phenolphthalein, the results being expressed in ml. of 0.01 N lye to 1 g. of abs. dry wheat. The moisture was detd. by noting the change in sp. gr. of H_2SO_4 in a desiccator in which samples of the grain were kept. Lipase activity and moisture were directly related. Protein was also detd. The wheats contg. more protein exhibit a lesser hygroscopicity. Even considerable differences in the chem. compn. of the wheat exercise only a very small influence on its moisture content (max. fluctuation 0.7%). The differences between wheat and the other cereals examd. are negligible. Nine references. A. H. K.

ASD-11A METALLURGICAL LITERATURE CLASSIFICATION

SECTION DIVISION

SECTION DIVISION

SECTION DIVISION

SECTION DIVISION

SECTION DIVISION

SECTION DIVISION

SECTION DIVISION

SECTION DIVISION

SECTION DIVISION

SECTION DIVISION

SECTION DIVISION

SECTION DIVISION

SECTION DIVISION

SECTION DIVISION

SECTION DIVISION

SECTION DIVISION

SECTION DIVISION

SECTION DIVISION

SECTION DIVISION

SECTION DIVISION

SECTION DIVISION

SECTION DIVISION

SECTION DIVISION

SECTION DIVISION

SECTION DIVISION

SECTION DIVISION

SECTION DIVISION

SECTION DIVISION

SECTION DIVISION

SECTION DIVISION

SECTION DIVISION

SECTION DIVISION

SECTION DIVISION

SECTION DIVISION

SECTION DIVISION

SECTION DIVISION

SECTION DIVISION

SECTION DIVISION

SECTION DIVISION

SECTION DIVISION

SECTION DIVISION

SECTION DIVISION

SECTION DIVISION

SECTION DIVISION

SECTION DIVISION

SECTION DIVISION

SECTION DIVISION

SECTION DIVISION

SECTION DIVISION

SECTION DIVISION

SECTION DIVISION

SECTION DIVISION

SECTION DIVISION

SECTION DIVISION

SECTION DIVISION

SECTION DIVISION

SECTION DIVISION

SECTION DIVISION

SECTION DIVISION

SECTION DIVISION

SECTION DIVISION

SECTION DIVISION

SECTION DIVISION

SECTION DIVISION

SECTION DIVISION

SECTION DIVISION

SECTION DIVISION

SECTION DIVISION

SECTION DIVISION

SECTION DIVISION

SECTION DIVISION

SECTION DIVISION

SECTION DIVISION

SECTION DIVISION

SECTION DIVISION

11A *No. 2* *11A*

Processes and Properties No. 11

Critical humidity and gas metabolism of stored Russian wheat and rye. V. L. Kretovich and E. N. Ushakov. *Compt. rend. acad. sci. U. R. S. S.* 29/115 19(1940) (in German). - Crit. humidity, detd. from the R. Q. by means of Sournov-Chigirev's apparatus at 25° was 15.5% for normal stored wheat and rye seed. Tests with CoCl_2 yielded the following values (in %) for "bound" water: filter paper 10.4; potato starch 12.4; wheat starch 11.8; gliadin 19.1. Since wheat and rye seeds consist largely of starch and protein, it is natural that the crit. humidity should lie between 15 and 10%, i. e., between the "bound" water content in protein and starch. With a low humidity a large part of the CO_2 eliminated is of anaerobic origin; only beyond the crit. humidity does the R. Q. approach the unit value. Other expts. showed a much more intensive respiration of germinated than of normal seeds of the same humidity. 11 references

A. H. Krapp

CA

114

Biochemical changes in wheat during its maturing after the harvest V. I. Kizilovskii and I. A. Akimovich (Doklady 6, No 4/6, 341-343 (1941) (German summary)).

The activity of the oxidation-reduction enzymes and amylase, the nonprotein N compounds, the acidity of extracts, and the properties of bran were detd. in freshly harvested, dried and after-matured wheat grains. Drying fresh grain in sun or with water ad diminishes the activity of the oxidation-reduction enzymes, the content of low mol. N compounds, and the alkali fixing substances. Drying increases the germinative capacity if the germination is done at 20°, but it lowers it if the germination is done at 10°. During the after-maturing, the synthesizing processes, which go on during the growth, terminate. This is expressed in a diminished content of nonprotein N compounds and of alkali-titratable substances in an extract of the grain. The elasticity of the bran decreases gradually. M. Horsch.

INSTITUTE OF BIOCHEMISTRY OF THE ACADEMY OF SCIENCES, USSR, MOSCOW

ca

2

Sören Peter Lauritz Sörensen. 1866-1940. V. L.
Kretovich. Uspokhi Khim. 10, 111-12(1941).
P. H. Rathmann

ASH-SLA METALLURGICAL LITERATURE CLASSIFICATION
RECEIVED MAR 1971

CA 12

Microbiological and biochemical processes in spontaneous heating of newly harvested wheat grain. V. L. Kretovich and Ya. I. Kautshitskiy. *Microbiology* (U. S. S. R.) 10, 401-8 (in English, 468) (1941); cf. C. A. 35, 4854. When freshly harvested wheat is piled in heaps on the threshing floor, changes in its microflora and in its biochem. properties set in at once. When the temp. inside a heap exceeds 35°, the germination capacity of the grain is lowered. This coincides with the first appearance of mold on the grain. By the next day profound autolytic processes are under way. The action of amylase and the acidity of alk. exs. increase, catalytic activity and gluten viscosity decrease. At 40° *B. mesentericus* and fungi develop rapidly, supplanting the specific microflora of sound grain. The increase in the no. of micrococci serves as an index of initial damage to the grain. On the 3rd day the micrococci are crowded out by fungi. Spontaneous heating of grain is a complicated process caused by the simultaneous action of enzymes and microorganisms. T. Laane

ASD-SLA METALLURGICAL LITERATURE CLASSIFICATION

10000 1000000000	100000 1000000000	1000000 1000000000	10000000 1000000000	100000000 1000000000	1000000000 1000000000
1000000000 1000000000	1000000000 1000000000	1000000000 1000000000	1000000000 1000000000	1000000000 1000000000	1000000000 1000000000

KRETOVICH, V. L.

"Concerning the Causes of the Reduced Germinative Capacity of Freshly-Harvested Corn,"
Dok. AN, 33, No. 2, 1941.

PROCESSES AND PROPERTIES INDEX

Respiration in seeds of flax and hulled cereals. V. I. Kretovskh. *Comp. rend. acad. sci. U. R. S. S.* 33:355-7 (1941).--Respiration in flax is greater than in hulled cereals of the same moisture content because of its high fat content. On the basis of the same moisture content in the hydrophilic portion of the seeds, the respiration rates are similar. The respiration coeff. is seldom 1.0. When moisture is low the evolution of CO_2 is greater than the absorption of O_2 , and is less at high moisture contents. The most precise method of detg. the loss of dry matter during storage is to det. it directly from the abs. dry wt.

J. T. Sullivan

ASB-SLA DETALLURGICAL LITERATURE CLASSIFICATION

62-2-4

NO. 5 1/D

110

CO

PROCESSES AND PROPERTIES

The dehydrogenases of wheat embryos. V. L. Kretovich and A. I. Sokolova. *Biokhimiya* 7, 232-7 (1942).— The losses attendant on grain storage over long periods of time or under unfavorable conditions are due to the weakening of the respiratory metabolism of the grain, as well as to the activity of the oxidation-reduction enzymes of the embryo, especially the dehydrogenases. In acid media, the wheat embryo dehydrogenases act very weakly and are completely inactive at pH 4.5-5. The optimum action for Mellin's buffer is at pH 7.2-7.5, and for Sørensen's phosphate buffer, pH 7.3-8.2. The optimum temp. is 50°. The dehydrogenase action is considerably enhanced in the presence of glutamic acid and hexose phosphates. H. Priestley

INST. OF BIOCHEMISTRY OF THE ACADEMY OF SCIENCES, USSR,

ASB 55A BIOLOGICAL LITERATURE CLASSIFICATION

1106

ca

1st and 2nd copies

PROCESSES AND PROPERTIES INDEX

120 and 4th copies

Common Element

Common Variable Index

Mechanism of wheat injury by *Eurygaster integriceps*.
V. L. Kretovich, A. A. Hundel and K. V. Pshenova.
Compt. rend. acad. sci. U. R. S. S. 39, 31-3 (1943).— No. 1
Enzymes of the salivary glands and anterior intestine dis-
sected from the wheat bug *Eurygaster integriceps* were in-
vestigated. During the period of wheat-stalk formation
the organs were found to contain a proteinase capable of
splitting casein, but not the gluten of normal wheat flour.
The salivary glands contained an amylase. During the
period of wax-ripening of the grain the salivary glands
contained an enzyme, extd. by dil. carbonate, which was
capable of destroying gluten. This enzyme exhibited
optimum activity at pH 8 and appeared to be a *trypsin*.
During this period the intestine contained sulphydryl
comps. capable of activating the protolytic enzymes of
normal wheat flour. R. E. Reeves

ASB-SLA METALLURGICAL LITERATURE CLASSIFICATION

FROM DIVISION

RECORD NO.

CLASSIFICATION

RECORD NO.

12

PROCESSED AND PREPARED BY

Proteolysis in grains affected with *Eurygaster integriceps*. V. I. Kretovich, K. V. Pukhova and A. A. Bun-
del. *Doklady Akad. Nauk S. S. R.* 40, 35 (1963),
Compt. rend. acad. sci. U. R. S. S. 40, 30 (1963) (in
English). --For tests for tyrosine, in free and peptide
form, in exts. from grain punctured by *Eurygaster integri-*
ceps (1) indicate that it secretes a proteinase of the trypsin
group. Samples of normal flour were mixed with aq. exts.
from grain damaged by 1, and with various glycerol buffer
solns. made up to different pH values. After standing at
20° for 2 hrs., the dough samples were tested for extensi-
bility of the gluten. Acidulation of the dough samples
inhibited protein splitting by the proteinase derived from
1. Max. protein splitting was observed at pH 6.6
I. W. HERS

with pure O. Low temps. favor germination because of higher O soly. in water and inhibition of molding. O may oxidize some inhibitors of germination or hasten enzyme formation in the embryo. During afterripening, there is produced first weaker, then stronger, gluten; this is correlated with a rise and fall of proteolysis. Acid-alc. extractives and low-mol. N compds. decrease as protein synthesis proceeds. Drying the grain decreases the reduction oxidation activity of its enzymes, the amylolytic activity and the content of physiologically unstable substances. Grain injured by shield-bugs (Pentatomidae) produces weak gluten. Glands of the salivary glands of the insects during their leaf-feeding stage are amylolytic and contain proteinase, which digests casein but not flour gluten. When the insects are grain-feeding, the glands contain amylase and trypsin, which digest gluten completely. The midgut contains thermostable glutathione-like sub-

stances that activate the enzymes in normal flour. The insect injury lowers the total N and gliadin of the grain; the gliadin becomes very soft, its viscosity decreases, and rotation changes, and S content increases. The proteolysis releases much free and peptidized tyrosine. Insect-induced proteolysis is most evident in soda extra; it is not due to cysteine-like activators. It is favored by neutral reaction and can be checked by acidification, adding AcOH bacteria or yeast liquid to the dough, dough formation at reduced temps, adding KI or H_2O_2 , or best by heating the grain, preferably with a quick steam treatment. Grain that is frozen before harvest produces flour of low H_2O -absorptive quality and poor baking properties; it is abnormally high in total and nonprotein N, diastatic power, acidity, and content of H_2O -sol. matter, and produces less elastic gluten, owing to proteolysis or protein coagulation; its gluten content is low, and its amylase shows strong saccharification and destruction. Vitreous grain is slightly more hygroscopic than mealy grain, but marked chemical differences in grain have little effect on its hygroscopic quality. The grain embryo contains more hygroscopic H_2O than the endosperm. The favorable effect of high temps. on H_2O absorption does not follow van't Hoff's law, because the process is biodynamic and not purely physical. Movement of H_2O to cool fuel in grain masses (unpublished data of Vasileva and Tsiganikova) is ascribed to thermosdiffusion and H_2O condensation, with the moisture gradient approx. proportional to the temp. gradient. In grain below 10% moisture content, much of the CO_2 is produced anaerobically, thus reduction-oxidation processes in stored grain are conditioned by moisture and not governed by laws of aerobic respiration. Enzymic processes

condition respiratory studies in indicating that the critical moisture of stored grain is 11.5-13.0%. In old, nonviable grain amylase is fully active. Changes in proteolytic activity do not follow viability changes. Titratable acidity increases with age of seed. There is close correlation between loss of dehydrase activity of embryos ($Na_2S_2O_3$ test) and of viability. Respiration falls with viability. Moistening embryos increases their dehydrase activity (methylene blue test), but later the enzyme is inactivated by molds. The max. activity is at pH 7.3-9.2 with little pH effect over a wide zone, and at 50° with a sharp decrease at 55°. The dehydrase system of wheat embryos is activated by glutamic acid, leucine di- and monophosphate, and phosphogluconic acid as H donors. The wheat esterase which affects triacetin is greatest in flour, less in bran, and least in embryos. In the wheat-lipase action on olive oil, embryo is most active, bran least. The moisture temp. ranges permitting safe grain storage are shown graphically. In afterripening "sweating," due to syntesis, is considered the basic cause of "dry" spontaneous heating and molding of grain. Con- blin-harvested grain contg. 13-16% moisture is best stored with forced ventilation systems. Wheat with 10-23% moisture is best treated by heating at 45°. Above this temp. the grain loses viability. Heating reduces the content of ale-sol. matter, the elasticity of the gluten, and the titratable acidity of the grain; this indicates hastened afterripening. Baking quality and loaf vol. are improved. Wheat of 14-15% moisture content retains its viability if stored at -5° to -15°, and moist (20%) wheat can be safely stored 3 mo. at -5°. 149 Russian and 244 non Russian references
K. Starr Chester

KRETOVICH, V. L.

"New Method of Extraction of the Free Fatty Acids from Oil," Biokhim., Vol. 10,
No. 2, 1945.

INST. OF BIOCHEMISTRY IM. A.N. BAKH OF THE ACADEMY OF SCIENCES, USSR,
MOSCOW

P.159.

BC

a-1

Processes and Properties: Index

Oil from millet rendered toxic by wintering in the field. V. L. Kretovitch and A. A. Bouda (*Biochimica*, 1945, 18, 218-224).— The toxic principle was studied by the inflammatory reaction produced on injection into the rabbit's skin. Toxic millet contains a large proportion of steryl kernels, and ethereal extracts of these are especially rich in the toxin. Extraction with 60% ethanol at 10° completely removes the principle, and such extracts have high acid, chromatographic analysis suggests that the toxic principle is associated with the unsaturated fatty acids, and is probably an oxidation product of these. It is not associated with aldehydes formation. If the oil from normal millet is treated with Na_2CO_3 , a toxic principle can be produced. D. H. S.

INST. OF BIOCHEMISTRY IM. A.N. BAKH OF THE ACADEMY OF SCIENCES, USSR, MOSCOW

ASB SLA RETAILING LITERATURE CLASSIFICATION

12

Biochemical properties of toxic millet. V. Kretovskiy,
N. Mamedov, Z. Shripina, and V. Shvetsov (Kurmukov
Inst. Cereal Research). *Nisbhemys* 10, 379-84 (1948).--
Grain is often found to be toxic as a food if it has lain all
winter in the field, covered with snow. Toxic millet differs
from normal grain in having a higher content of nonpro-
tein and amino N, and a lower activity of mahlizing en-
zymes. Dextrin formation by amylase, as detd. by
Wohlgemuth's method, is twice as high in toxic millet.
H Priestley

INST. OF BIOCHEMISTRY IM.A.N. BAKH OF THE ACADEMY OF SCIENCES, USSR,
AND THE ALL-UNION GRAIN RESEARCH INSTITUTE

ASH-55A METALLURGICAL LITERATURE CLASSIFICATION

CIA-RDP86-00513R0008264200

<p>CA</p> <p>No. 2</p>		<p>PROCESSES AND PROPERTIES</p> <p>Inhibition of alcoholic fermentation by fat decomposition products. V. I. Kertovich, E. N. Mishustin, and A. A. Hundel (Bach Biochem. Inst., Moscow). <i>Biochimiya</i> 11, 149-54(1946); cf. C.A. 39, 8349. -Alc. fermentation is inhibited by the oil extd. from "toxic" millet, wheat, and rye (grain which had lain in the field all winter and which causes "septic angina"). Fat decomposition products are responsible for this powerful physiol. effect. The oil from grain affected by molds did not inhibit alcoholic fermentation. H. Priestley</p>	
<p>INST. OF BIOCHEMISTRY, ACADEMY OF SCIENCES IM. A.N. BAKH AND INST. OF MICROBIOLOGY OF THE ACADEMY OF SCIENCES, USSR, MOSCOW</p>		<p>ASSOCIATE METALLURGICAL LITERATURE CLASSIFICATION</p>	
<p>SHOW EXTENSIVE</p>		<p>SHOW BREVITY</p>	
<p>100000 01</p>		<p>011111 011</p>	

6.1

17

No. 6

V. E. Murty
 Substances producing the odor and bitter taste of grain contaminated with absinth. V. L. Kretovich, Yu. S. Rall, and L. A. Trisvyatskii (Inst. Food Technol., Moscow). *Biokhimiya* 11, 403-404 (1946).--In certain sections of Russia, the grain is contaminated with the dust of absinth (*Artemisia absinthium* and *A. hermodactyla*), which confer on the flour, and even the bread, a specific absinth odor (caused by essential oils) and a strong bitter taste (caused by the glycoside absinthin). The bitter taste is completely removed by washing the grain with luke-warm water. H. Priestley

INST. OF FOOD TECHNOLOGY, MOSCOW

ASB. SLA METALLURGICAL LITERATURE CLASSIFICATION

62

12

4

1st AND 2nd DEPT'S

PROCESSING AND PROPERTIES UNIT

FERMENTATION test as a diagnostic method for toxic hibernated grain. R. N. Mishustin, V. L. Kretovich, and A. A. Bundel. *Gigiena i Sanit.* 11, No. 11, 32-5 (1946).—The toxic principle which is present in grain which had spent the winter in the field sharply reduces the process of yeast growth and thus inhibits ale fermentation. This permits the use of a fermentation test for such grain, which is capable of producing alimentary alkalis. Tests (in lab. only) confirmed this method on wheat, rye, and millet. The KOH ext. of the grain (0.1-0.2% by wt.) is introduced into the yeast culture (0.5 million cells per ml.) in physiol. soln., the nutrient medium being most conveniently sugar; the tubes are incubated 2 days at 20° and the vol. of evolved gas is measured.

G. M. Kozelapoff

ASM-11A METALLURGICAL LITERATURE CLASSIFICATION

FROM SOURCE

REMARKS ON CHG. 111

11 12 13 14 15 16 17 18 19 20 21 22 23 24 25 26 27 28 29 30 31 32 33 34 35 36 37 38 39 40 41 42 43 44 45 46 47 48 49 50 51 52 53 54 55 56 57 58 59 60 61 62 63 64 65 66 67 68 69 70 71 72 73 74 75 76 77 78 79 80 81 82 83 84 85 86 87 88 89 90 91 92 93 94 95 96 97 98 99 100

KRETovich, V. L.

MISHUSTIN, E. N., KRETovich, V. L., and BUNDEL', A. A. "Fermentative Test as a Method of Diagnosing Toxicity of Grain," in Reports of the Scientific-Research Work for 1945, Department of Biological Science, Publishing House of the Academy of Science USSR, Moscow, 1947, p. 150. 511 Ak144

Sira-Si-90-53, 15 Dec. 1953

11-D

CA

Biochemistry of the ripening of rye grain. V. Kretovich, R. Tokareva, I. Petrova, T. Drozdova, L. Auerman, and N. Senolina (Baking Inst., Moscow). *Biochemistry* 12, 646-66(1947); cf. C.A. 30, 82619. As the grain ripens, there is a decrease in the amylase and proteolytic activity, and in the nonprotein N and reducing sugars. The farther the ripening proceeds, the less the proteins and starches are liable to be attacked by the enzymes. Early harvesting of the grain is not the cause of poor bread-baking qualities, but the slightest sprouting is highly detrimental. H. Priestley

ASB-SLA METALLURGICAL LITERATURE CLASSIFICATION

KRISTOVICH, V. L.

(fungal)

"Investigation of qumies of Rye Grain," Biokhim., Vol. 12, No. 12, 1947.

KRETovich, V., TOKAREVA, R., AUERMAN L., SMOLINA N., KULMAN A., AND BRANOPOL'SKAYA R.

"Change in the Quality of Rye Flour During Storage, " Dok. An, 58, No.9, 1947.

[illegible]

1ST AND 2ND ORDERS																										3RD AND 4TH ORDERS																									
PROCESSING AND PROPERTIES INDEX																										1ST AND 2ND ORDERS																									
100																										110																									
120																										130																									
140																										150																									
160																										170																									
180																										190																									
200																										210																									
220																										230																									
240																										250																									
260																										270																									
280																										290																									
300																										310																									
320																										330																									
340																										350																									
360																										370																									
380																										390																									
400																										410																									
420																										430																									
440																										450																									
460																										470																									
480																										490																									
500																										510																									
520																										530																									
540																										550																									
560																										570																									
580																										590																									
600																										610																									
620																										630																									
640																										650																									
660																										670																									
680																										690																									
700																										710																									
720																										730																									
740																										750																									
760																										770																									
780																										790																									
800																										810																									
820																										830																									
840																										850																									
860																										870																									
880																										890																									
900																										910																									
920																										930																									
940																										950																									
960																										970																									
980																										990																									
1000																										1010																									
1020																										1030																									
1040																										1050																									
1060																										1070																									
1080																										1090																									
1100																										1110																									
1120																										1130																									
1140																										1150																									
1160																										1170																									
1180																										1190																									
1200																										1210																									
1220																										1230																									
1240																										1250																									
1260																										1270																									
1280																										1290																									
1300																										1310																									
1320																										1330																									
1340																										1350																									
1360																										1370																									
1380																										1390																									
1400																										1410																									
1420																										1430																									
1440																										1450																									
1460																										1470																									
1480																										1490																									
1500																										1510																									
1520																										1530																									
1540																										1550																									
1560																										1570																									
1580																										1590																									
1600																										1610																									
1620																										1630																									
1640																										1650																									
1660																										1670																									
1680																										1690																									
1700																										1710																									
1720																										1730																									
1740																										1750																									
1760																										1770																									
1780																										1790																									
1800																										1810																									
1820																										1830																									
1840																										1850																									
1860																										1870																									
1880																										1890																									
1900																										1910																									
1920																										1930																									
1940																										1950																									
1960																										1970																									
1980																										1990																									
2000																										2010																									
2020																										2030																									
2040																										2050																									
2060																										2070																									
2080																										2090																									
2100																										2110																									
2120																										2130																									
2140																										2150																									
2160																										2170																									
2180																										2190																									
2200																										2210																									
2220																										2230																									
2240																										2250																									
2260																										2270																									
2280																										2290																									
2300																										2310																									
2320																										2330																									
2340																										2350																									
2360																										2370																									
2380																										2390																									
2400																										2410																									
2420																										2430																									
2440																										2450																									
2460																										2470																									
2480																										2490																									
2500																										2510																									
2520																										2530																									
2540																										2550																									
2560																										2570																									
2580																										2590																									
2600																										2610																									
2620																										2630																									
2640																										2650																									
2660																										2670																									
2680																										2690																									
2700																										2710																									
2720																										2730																									
2740																										2750																									
2760																										2770																									
2780																										2790																									
2800																										2810																									
2820																										2830																									
2840																										2850																									
2860																										2870																									
2880																										2890																									
2900																										2910																									
2920																										2930																									
2940																										2950																									
2960																										2970																									
2980																										2990																									
3000																										3010																									
3020																										3030																									
3040																										3050																									
3060																										3070																									
3080																										3090																									
3100																										3110																									
3120																										3130																									
3140																										3150																									
3160																										3170																									
3180																										3190																									
3200																										3210																									
3220																										3230																									
3240																										3250																									
3260																										3270																									
3280																										3290																									
3300																										3310																									
3320																										3330																									
3340																										3350																									
3360																										3370																									
3380																										3390																									
3400																										3410																									
3420																										3430																									
3440																										3450																									
3460																										3470																									
3480																										3490																									
3500																										3510																									
3520																										3530																									
3540																										3550																									
3560																										3570																									
3580																										3590																									
3600																										3610																									
3620																										3630																									
3640																										3650																									
3660																										3670																									
3680																										3690																									
3700																										3710																									
3720																										3730																									
3740																										3750																									
3760																										3770																									
3780																										3790																									
3800																										3810																									
3820																										3830																									
3840																										3850																									
3860																										3870																									
3880																										3890																									
3900																										3910																									
3920																										3930																									
3940																										3950																									
3960																										3970																									
3980																										3990																									
4000																										4010																									
4020																										4030																									
4040																										4050																									
4060																										4070																									
4080																										4090																									
4100																										4110																									
4120																										4130																									
4140																										4150																									
4160																										4170																									
4180																										4190																									
4200																										4210																									
4220																										4230																									
4240																										4250																									
4260																										4270																									
4280																										4290																									
4300																										4310																									
4320																										4330																									
4340																										4350																									
4360																										4370																									
4380																										4390																									
4400																										4410																									
4420																										4430																									
4440																										4450																									
4460																										4470																									
4480																										4490																									
4500																										4510																									
4520																										4530																									
4540																										4550																									
4560																										4570																									
4580																										4590																									
4600																										4610																									
4620																										4630																									
4640																										4650																									
4660																										4670																									
4680																										4690																									
4700																										4710																									
4720																										4730																									
4740																										4750																									
4760																										4770																									
4780																										4790																									
4800																										4810																									
4820																										4830																									
4840																										4850																									
4860																										4870																									
4880																										4890																									
4900																										4910																									
4920																										4930																									
4940																										4950																									
4960																										4970																									
4980																										4990																									
5000																										5010																									
5020																										5030																									
5040																										5050																									
5060																										5070																									
5080																										5090																									
5100																										5110																									
5120																										5130																									
5140																										5150																									
5160																										5170																									
5180																										5190																									
5200																										5210																									
5220																										5230																									
5240																										5250																									
5260																										5270																									
5280																										5290																									
5300																										5310																									
5320																										5330																									
5340																										5350																									
5360																										5370																									
5380																										5390																									
5400																										5410																									
5420																										5430																									
5440																										5450																									
5460																										5470																									
5480																										5490																									
5500																										5510																									
5520																										5530																									
5540																										5550																									
5560																										5570																									
5580																										5590																									
5600																										5610																									
5620																										5630																									
5640																										5650																									
5660																										5670																									
5680																										5690																									
5700																										5710																									
5720																										5730																									
5740																										5750																									
5760																										5770																									
5780																										5790																									
5800																										5810																									
5820																										5830																									
5840																										5850																									
5860																										5870																									
5880																										5890																									
5900																										5910																									
5920																										5930																									
5940																										5950																									
5960																										5970																									
5980																										5990																									
6000																										6010																									
6020																										6030																									
6040																										6050																									
6060																										6070																									
6080																										6090																									
6100																										6110																									
6120																										6130																									
6140																										6150																									
6160																										6170																									
6180																										6190																									
6200																										6210																									
6220																										6230																									
6240																										6250																									
6260																										6270																									
6280																										6290																									
6300																										6310																									
6320																										6330																									
6340																										6350																									
6360																										6370																									
6380																										6390																									
6400																										6410																									
6420																										6430																									
6440																										6450																									
6460																										6470																									
6480																										6490																									
6500																										6510																									
6520																										6530																									
6540																										6550																									
6560																										6570																									
6580																										6590																									
6600																										6610																									
6620																										6630																									
6640																										6650																									
6660																										6670																									
6680																										6690																									
6700																										6710																									
6720																										6730																									
6740																										6750																									
6760																										6770																									
6780																										6790																									
6800																										6810																									
6820																										6830																									
6840																										6850																									
6860																										6870																									
6880																										6890																									
6900																										6910																									
6920																										6930																									
6940																										6950																									
6960																										6970																									
6980																										6990																									
7000																										7010																									
7020																										7030																									
7040																										7050																									
7060																										7070																									
7080																																																			

KRETOVICH, V. I.

USSR/Medicine -Plants
Medicine -Metabolism

Mar/Apr 48

"Problem of Metabolism of Plants at the Fourth All-Union Botanical Convention,"
A. I. Oparin, V. I. Kretovich, 1 $\frac{1}{2}$ pp

"Botan Zhur" No 2

PA 28/1:2764

Dmitrii Nikolaevich Pryanishnikov (1865-1948): A. I.
 Knyazevskiy (Nach. Bushen Inst., Moscow) 16.
 Akademiya 13, 283-5 (1948). An obituary of the foremost
 Russian plant physiologist and agricultural chemist.
 H. Preshev

KRETOVICH, V. L.

PA 12/49T1

USSR/Academy of Sciences
Biography

Jul/Aug 1948

"In Memory of Academician Dimitriy Nikolayevich
Pryanishnikov," V. L. Kretovich, 2 3/4 pp

"Biokhimiya" Vol XIII, No 4

Obituary notice of D. N. Pryanishnikov, eminent
agrochemist, physiologist and biologist. Photograph
shows head and shoulders (CIA Photo Accession No
3108).

12/49T1

CA

No. 4

Sulfhydryl compounds and ascorbic acid in sprouting and ripening seeds. V. L. Kretovich, A. A. Bundel, and T. V. Prozdova (Bach Biochem. Inst., Moscow). *Plants* 13, 432-6 (1945).--The glutathione and ascorbic acid contents in seeds (wheat, rye, corn) rise sharply at the very beginning of germination, reach a definite max., and then decline. The concn. of sulfhydryl compds. is particularly high in the leaf and root; it is less in the cyme, and least in the endosperm. Ascorbic acid is completely absent both in the cyme and in the endosperm. H. Priestley

ASB-SLA METALLURGICAL LITERATURE CLASSIFICATION

CA

№ 6

Interaction of amino acids and sugars at high temperatures. V. L. Kretovich and R. Tokareva. *Biokhimiya* 13, 508-15(1948).—Melanoidin formation was measured by heating at 95° a soln. of 2 ml. distd. H₂O, 200 mg. sugar, and 60 mg. amino acid. The color caused by the melanoidins was compared with the color of standard I solns. The most intense melanoidin formation occurred in the presence of pentoses. Of the disaccharides, maltose reacted, but sucrose did not; hence a free carboxyl group is necessary for melanoidin formation. Glycine was the most reactive of the amino acids. Then followed leucine, alanine, and other amino acids, including di- and tripeptides. Melanoidin formation was accompanied by the formation of furfural and other volatile aldehydes, which imparted the aroma to the meat. When the aldehydes were tied up by the addn. of dimedone, melanoidin formation was prevented. 11 Priestley

Inst. Biochem. im. A.-N. Bakh.

ASAC-114 DETAILED LITERATURE CLASSIFICATION

CA

Transformation of alime (soluble pentosans) during germination and ripening of rye seed. V. I. Kiselevich and I. S. Petrova. Doklady Akad. Nauk S.S.S.R. 59, 281 3(1948). Rye seed, in contrast to other grains, has a considerable amt. of "alime" which represents the sol. pentosans, the transformations of which during growth have not been examd. previously. The present studies were made over 2 growing seasons. After 4 hrs. wetting the seeds were grown on wet filter paper. The changes in pentosans were followed by total detn. and estn. of high- and low-mol. wt. fractions (the former are pptd. by strong KOH, 70%, and higher). The ground grain was placed

with water in a graduated vessel, agitated, let stand 0.5 hr., centrifuged, and two 50-ml. aliquots were taken. The 1st was used for total-pentosan detn.; the 2nd aliquot was treated with 6 vols. 90% KOH and allowed to stand overnight; after filtration of the high-mol.-wt. fraction, the filtrate was again checked for total pentosan content. Control of pentose content was made by extrn. of ground grain with 80% alc. at 75°, followed by evapn., soln. in H₂O, and fermentation by bakers' yeast conventionally. Depending on the location from which samples were taken, the total pentosans ranged from 2.47 to 1.34% (on dry wt.) during ripening, and rose to 3.1-4.07% during germination (3rd and 6th day, resp.). The high-mol. pentosans remained at 1.8-1.9% level during early ripening, dropped to 1.3 at ripeness and rose to 2.76-2.98% during germination. The low-mol.-wt. pentosans dropped from 0.67 to 0.01% in ripening and rose to 0.37-1.09% in germination (3rd and 6th day, resp.). The ripening process covered approx. 2 months. G. M. Kosolapoff

Inst. Bread Baking Industry, Moscow

PA 51145

KRETOVICH, V. L.

21 Mar 1948


USSR/Medicine - Plants
Medicine - Alanine

"Synthesis of Alanine in Vegetable Tissues," V. L.
Kretovich, A. A. Bundel', Inst Biochem imeni A. N.
Bakh, Acad Sci USSR, 4 pp

"Dok Akad Nauk SSSR, Nova Ser" Vol LIX, No 9

Reports experimental research on the conditions of
synthesis of alanine from pyruvate in ground and
living tissues of plants. Experiments conducted
with lupine and squash, chosen as characteristically
representing two types of oxygen exchange in plants.
Presents process of experiments and results. Sub-
mitted by Academician A. I. Operin, 24 Jan 1948.

51T45



[illegible]

110

CA

Oxidation of amino acids by plant tissues. V. Kretovich and T. Drozlova. *Doklady Akad. Nauk S.S.S.R.* 63, 167-70 (1948). - Manometric studies, with oxidative enzymes from rye sprouts (phosphate buffer), with amino acid substrate at 31° showed: O uptake is sharply increased by pH change from 5 to 8 without the enzymes, this effect is absent in the complete system and sharpest results are observed at pH 5, which was then used. In carboxylic amino acids (aspartic and glutamic) give 3.2 times greater O uptake than the monocarboxylic acids. Lysine, tryptophan, and glycine are but little attacked. Tyrosine, alanine, proline, cystine, leucine and arginine are nearly identical and are oxidized 3-4 times more energetically than lysine; histidine is twice as active as alanine. Glutamine not only is not oxidized, but suppresses oxidation of normal substrates. Oxidation of glutamic acid is almost completely inhibited by 0.01 M HCN.

G. M. Kowaloff

KRETOVICH, V. L. i YEVSTIINEYEVA, Z. G.

20014 KRETOVICH, V. L. i YEVSTIINEYEVA, Z. G. O nakhozhdanii glyutamina v sakharney
svekle. Biokhimiya, 1949, Vyp. 3, s. 271-74. -- Bibliogr: 7 nazv.

SO: LETOPIS ZHURNAL STATEY, Vol. 27, Moskva, 1949.

CA


112


Presence of glutamine in the sugar beet V. I. Kretovich and Z. G. Ivastigneeva. *Biokhimiya* 14, 271 (1949). Contrary to the opinion expressed in the literature, the Russian sugar beet does contain glutamine. There is only a very small amt. of asparagine and glutamine in the leaves of the sugar beet throughout the various stages of its development. In the roots, the amt. of the amides increases, reaching a max. around Sept., when the asparagine and glutamine N is 0.12% on a dry basis, or 11% of the total N of the root. H. Priestley

Inst. Biochem. in A.N. Bakh, AS USSR

ASB SLA DETAILING LITERATURE CLASSIFICATION

1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28	29	30	31	32	33	34	35	36	37	38	39	40	41	42	43	44	45	46	47	48	49	50	51	52	53	54	55	56	57	58	59	60	61	62	63	64	65	66	67	68	69	70	71	72	73	74	75	76	77	78	79	80	81	82	83	84	85	86	87	88	89	90	91	92	93	94	95	96	97	98	99	100
---	---	---	---	---	---	---	---	---	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	-----





CA

12

Volatile aromatic constituents of bread and malt. V. I. Krasavich and R. R. Tokareva (Vsesoyuz. Nauch.-Issledovatel. Inst. Khlebopekarnoi Prom., Ministerstva S.S.S.R. 66, 231-4 (1949).—Steam distn. in H_2O at 40°C showed that the pleasant taste and olfactory character-istics of bread of red rye malt are paralleled by volatile aldehyde content. Generally, 21-30 mg./100 g. (calcd. as AcH) can be regarded as the dividing line for malt-classification. In bread, the "higher" types of wheat bread are lower in aldehydes than the more "crude" forms of dark bread, which are more aromatic; these values range from 3 to 9 mg./100 g. as AcH, with essential absence of furfural in refined types of wheat bread, ranging to 0.8 in rough dark rye bread. Traces of acetylmethylcarbinol are found in malt, but not in the bread. In addition, volatile acids and esters also contribute to the aroma of the materials. The identification of individuals is not accomplished. (1) M. Krasavich

all-Union Sci. Res. Inst. Bread-Baking Industry, Mining Food, USSR

12

CA

Dependence of grain respiration on temperature.
A. P. Frokhorova and V. L. Kretovich (Research Inst. of
Ministry of Material and Agr. Reserves, and A. N. Bakh
Biochem. Inst., Moscow). Doklady Akad. Nauk S.S.
S.R. 69, 401-3 (1949).—Optimum respiration temp. for
wheat grain is 30-35° and the temp. coeff. (10° interval)
varies: for 0-10° it is 5 at 14% moisture content, 22 at
16%, 7.2 at 18%, and 12.0 at 22% moisture; for 10-20°
intervals these are 8, 2.9, 6.2, and 3.6; for 20-30° inter-
vals: 2.7, 2.4, 2.7, 2.1; for 30-40°: 2.3, 2.2, 2.2, 2.9;
for 40-50°: 2.0, 1.6, 1.7, 1.6, resp. Grain with 14-16%
moisture keeps const. respiration rate at 35° for several
days, but on long exposures the rate declines, while grains
with high moisture level (18-23%) begin to decline even
after 6-18 hrs. (G. M. Kowaloff)

(BA - A III Mr. 58:395)

KRETOVICH, V. L., jt. au.

Koz'mina, N. P. Biochemistry of grain and products obtained from processing it;
textbook 4. ispr. i dop. izd. Moskva, Gos. izd-vo tekhn. i ekon. Lit-ry po voprosam
zagotovok, 1950. 358 p. (55-40820) Biokhimiia zerna...1950 (Card 2, 55-40820)

QK861.k615 1950

KRETovich, V. L.

Chem ③

Determination of aspartic and glutamic acids by the method of chromatographic adsorption. V. L. Kretovich and A. A. Buzilev. *Trudy Vsesoyuz. Sovetskaniya Khromatog., Akad. Nauk S.S.S.R., Otdel. Khim. Nauk* 1950, 192-9 (Pub. 1952).-- Sepn. of aspartic and glutamic acids on Al_2O_3 is described in detail. The latter is completely eluted by 0.5N AcOH, while the former is but slightly shifted downward during such elution. After this sepn. the aspartic acid can be removed by elution with alkali. The analysis consists of washing the adsorbent with 6N HCl, followed by H_2O until the pH reaches 2.5-2.7, after which the biol. soln. is sucked through the adsorbent tube, washed with H_2O satd. with H_2S , then plain H_2O , eluted with 0.5N AcOH, distd. H_2O , then with 3N and finally 0.05N KOH. The solns. are collected separately and are used for the usual combustion method of N detn., from which the content of the acids is calcd. In young leaves and stems of wheat both aspartic and glutamic acids are present in equal amts. Sepn. of 10 mg. is possible, with an error of about 5%. G. M. K.

7

Chromatographic separate determination of aspartic and glutamic acids. V. L. Kirilovskiy and A. A. Bunde (A. N. Bakht. Khim. Ind. Acad. Sci. U.S.S.R.) Doklady Akad. Nauk S.S.S.R. 73, 117 (1950). Samples (1 g.) are treated with hot 10% KOH, after which the treatment is as given earlier (C.I. 43, 3745). The anionotropic Al_2O_3 , however, is prepd. from 10 g Al_2O_3 which is treated with 30 ml. 6 N HCl, then washed to pH 2.5-2.7. The final product is heated with almost as much $AlCl_3$ for 24 hrs. at 70°C to give the most active product. Adsorption is done in a (6) cm. x 8 x 10 cm. tube at 16-18°, by passing 2-4 ml. ext. with suction through the packed tube, washing with 50 ml. H_2O and by H_2S , then H_2O , followed by elution of glutamic acid by 55 ml. 0.5 N AcOH, then H_2O , and elution of aspartic acid by 5 ml. 3 N KOH then 40 ml. 0.05 N KOH. The separate eluates are then decompd. as usual and the microdetn. of N is performed. Etiolated lupine sprouts were found to contain 25.75% aspartic and 21% glutamic acids. wheat (leaves and stem): 24.40 and 33.6% resp. Separation of 10 mg. of each acid is perfectly feasible by this method. G. M. Kosolapoff.